

## Research article

---

# A simple clinical coding strategy to improve recording of child maltreatment concerns: an audit study

Cite this article: McGovern A, Woodman J, Allister J, van Vlymen J, Liyanage H, Jones S, Rafi I, de Lusignan S, Gilbert R. A simple clinical coding strategy to improve recording of child maltreatment concerns: an audit study. *Inform Prim Care*. 2014;22(1):227–234.

<http://dx.doi.org/10.14236/jhi.v22i1.93>

Copyright © 2014 The Author(s). Published by BCS, The Chartered Institute for IT under Creative Commons license <http://creativecommons.org/licenses/by/4.0/>

**Author address for correspondence:**

Simon de Lusignan  
Clinical Informatics,  
University of Surrey,  
Guildford GU2 7PX, UK  
Email: [s.lusignan@surrey.ac.uk](mailto:s.lusignan@surrey.ac.uk)

Accepted October 2014

**Andrew McGovern**

Clinical Informatics, Department of Health Care Management and Policy, University of Surrey, Guildford, UK

**Jenny Woodman**

Population, Policy and Practice, Institute of Child Health, University College London, London, UK

**Janice Allister**

Royal College of General Practitioners, London, UK

**Jeremy van Vlymen**

Clinical Informatics, Department of Health Care Management and Policy, University of Surrey, Guildford, UK

**Harshana Liyanage**

Clinical Informatics, Department of Health Care Management and Policy, University of Surrey, Guildford, UK

**Simon Jones**

Clinical Informatics, Department of Health Care Management and Policy, University of Surrey, Guildford, UK

**Imran Rafi**

Royal College of General Practitioners, London, UK

**Simon de Lusignan**

Clinical Informatics, University of Surrey, Guildford, UK

**Ruth Gilbert**

Population, Policy and Practice, Institute of Child Health, University College London, London, UK

---

## ABSTRACT

**Background** Recording concerns about child maltreatment, including minor concerns, is recommended by the General Medical Council (GMC) and National Institute for Health and Clinical Excellence (NICE) but there is evidence of substantial under-recording.

**Aim** To determine whether a simple coding strategy improved recording of maltreatment-related concerns in electronic primary care records.

**Design and Setting** Clinical audit of rates of maltreatment-related coding before January 2010–December 2011 and after January–December 2012 implementation of a simple coding strategy in 11 English family practices. The strategy included encouraging general practitioners to use, *always and as a minimum*, the Read code ‘Child is cause for concern’. A total of 25,106 children aged 0–18 years were registered with these practices. We also undertook a qualitative service evaluation to investigate barriers to recording.

**Method** Outcomes were recording of 1) any maltreatment-related codes, 2) child protection proceedings and 3) child was a cause for concern.

**Results** We found increased recording of any maltreatment-related code (rate ratio 1.4; 95% CI 1.1–1.6), child protection procedures (RR 1.4; 95% CI 1.1–1.6)

and cause for concern (RR 2.5; 95% CI 1.8–3.4) after implementation of the coding strategy. Clinicians cited the simplicity of the coding strategy as the most important factor assisting implementation.

**Conclusion** This simple coding strategy improved clinician's recording of maltreatment-related concerns in a small sample of practices with some 'buy-in'. Further research should investigate how recording can best support the doctor–patient relationship.

**Keywords:** child abuse, clinical audit, clinical coding, informatics, interdisciplinary communication

**How this fits in** Recording concerns about child maltreatment, including minor concerns, is recommended by the General Medical Council (GMC) and National Institute for Health and Clinical Excellence (NICE), but there is evidence of substantial under-recording. We describe a simple clinical coding strategy that helped general practitioners to improve recording of maltreatment-related concerns. These improvements could improve case finding of children at risk and information sharing.

## INTRODUCTION

Child safeguarding (Box 1) includes the recording of concerns and information about vulnerable children in the child's medical record.<sup>1</sup> There is substantial under-recording of child maltreatment and maltreatment-related concerns in primary care records,<sup>2,3</sup> which places children at increased risk of harm.<sup>4–6</sup> Recording of concerns underpins sharing of information between practitioners, failure of which is highlighted repeatedly as an important factor contributing to serious adverse outcomes.<sup>7</sup> The NICE<sup>8</sup> and the GMC<sup>9</sup> recognise that there is a spectrum of concerns that includes minor concerns. Their guidelines emphasise recording the whole spectrum of concerns. However, clinicians are apprehensive about how recording is perceived by parents and children and the impact of this on the patient–doctor relationship.<sup>3</sup>

### Box 1 Definition of child safeguarding

In England, child safeguarding is defined by statutory guidance<sup>10</sup> as comprising:

- protecting children from maltreatment;
- preventing impairment of children's health or development;
- ensuring that children grow up in circumstances consistent with the provision of safe and effective care, and
- taking action to enable all children to have the best outcomes.

England has a registration-based system for primary care with patients permanently registered with just one practice. Primary care notes are universally computerised with records made at the time a patient is seen,<sup>11</sup> although other information from test results and letters is often added to the computerised record outside the consultation. Data entered into the patient record are either coded (a process of assigning a clinical code to the patient record with associated values) or entered as free-text. In current clinical systems only coded

data are searchable. Therefore, careful clinical coding of maltreatment concerns is essential to allow a cumulative picture of concerns to emerge and for any kind of proactive management of patients who prompt concern. Good record keeping is also key for effective sharing of child maltreatment concerns.<sup>12</sup>

With the aim of improving coding of maltreatment-related concerns in general practice as recommended in NICE guidance,<sup>8</sup> we developed a simple coding strategy (Figure 1) using consensus methods with general practitioners (GPs) from 11 practices in England.<sup>3</sup> Here, we report the results of an audit comparing the rate of coding of concerns before and after implementation of the coding strategy in these practices. We also conducted a process evaluation that elicited feedback from practitioners on the potential barriers and facilitators for implementing the change in recording practice.

## METHODS

The characteristics of the 11 English practices instigating the intervention have been reported previously;<sup>3</sup> importantly, selection of these practices was based on identifying a 'GP lead' with known interest in either child protection or other relevant expertise, who could lead the implementation in their primary care practice.

### Implementing the coding strategy

The consensus developed coding strategy was implemented in January 2012 (Figure 2). Participating practices were sent implementation packs in mid-November 2011 to enable pre-implementation training and dissemination to members of the practice by GP leads. The implementation pack contained a letter summarising the study progress, a feedback form, a short presentation explaining the coding strategy and detailing the baseline rate of recording maltreatment-related codes.

As part of the coding strategy GPs were asked to use, *always and as a minimum*, the 'Child is cause for concern' Read code to document all maltreatment related concerns.




Figure 1 The simple coding strategy implemented. The solid lines represent the recommended minimum code use. Figure adapted from Ref.<sup>3</sup>




Figure 2 Flowchart diagram of the simple coding intervention and data collection methods.

## Measuring rates of coded maltreatment-related concern

We calculated the average rate of coded maltreatment-related concerns per year before implementation in the 11 practices, based on rates in each of two years (January 2010–December 2011). The ‘before implementation’ rates were compared with the rate of coded concerns in the 12 months following implementation (January–December 2012).

In England, most primary care data are recorded using the hierarchical Read coding system version 2, though a substantial minority of practices use read Clinical Terms version 3.<sup>13</sup> For our analysis we mapped all the codes to their read two equivalent (mappings available from <http://www.clininf.eu/maltreatment/>). A total of 350 relevant read codes were identified for extraction and analysis. The complete code list is provided in supplementary file 1 (SF1). The three outcome measures were recording as follows.

1. Any maltreatment-related code including the subgroups described below; a total of 350 relevant Read codes [SF1 and Table 1(a) and (b)].
2. Any child protection procedure; a total of 24 codes [SF 1 and Table 2(a) and (b)].
3. Any cause for concern code (SF 1 and Table 3); 1 code entitled ‘Child is cause for concern’ (SF1 and Table 3).

All patients under 18 who were permanently registered at the practices at the date of download in Dec 2012 were included in the analysis, even if they had not been registered throughout the whole period of the study. We excluded children who had been registered previously but had left the practice before the date of data download. Relevant anonymised recorded data (maltreatment-related codes, patient gender; registration date with the practice and patient age in years) were extracted using Morbidity Information Query and Export Syntax (MIQUEST).

We calculated the rate of maltreatment-related code recording per 1000 person years of risk, and an adjusted rate standardised for age and gender with 95% confidence intervals. A child could only be counted once in the numerator for each calendar year. Coding rates were, therefore, defined as the number of children with one or more maltreatment-related codes recorded per 1000 person years at risk in one calendar year. We took the mean rate for 2010 and 2011 as the rate for the period before the intervention.

We calculated adjusted rate ratios based on the rate after implementation divided by the rate before implementation. Where rate ratio 95% confidence intervals do not cross parity, we assume the rate difference to be statistically significant.

## Process evaluation

Understanding the extent to which practices were exposed to the coding strategy and the extent to which the audit ran as intended (programme fidelity) are important elements of clinical audit and evaluation. Feedback on the approach used to implement the change was collected from GP leads via a teleconference in Feb 2012 and a questionnaire sent out by email in July 2012.

## RESULTS

Of the original 11 practices invited to participate one practice was excluded from the final analysis because there was no lead to implement the coding strategy after the GP lead left the practice. The included practices were geographically spread across England, but there was a majority in the major population centres of London and the South East (Table 1). The complete characteristics of these practices have been described in detail elsewhere.<sup>3</sup> The included practices used a range of primary care software: three practices used EMIS LV; one practice EMISWEB, three practices INPS Vision, one practice Isoft Synergy and two practices TPP SystmOne.

**Table 1** Details of the primary care practices included in the study. IMD = index of multiple deprivation

Practice code	Region	IMD score (2010)	IMD decile	Audit lead remained	Included for analysis
P1	North East	47.6	1st	Yes	Yes
P2	Midlands	19.5	5th	Yes	Yes
P3	East	25.5	4th	No	Yes
P4	London	23.7	4th	Yes	Yes
P5	South East	39.3	2nd	Yes	Yes
P6	South East	3.61	10th	Yes	Yes
P7	South East	7.3	9th	Yes	Yes
P8	South East	4.42	10th	Yes	Yes
P9	South East	11.6	7th	Yes	Yes
P10	London	29.4	3rd	No	No
P11	London	19.3	5th	Yes	Yes

A total of 25,106 children were included; 20,975 children (36,140 person years) before the intervention and 21,783 children (19,560 person years) after the intervention. The age–gender profile of the population of children included were similar for the before and after intervention periods.

### Quantitative findings

During the two years before the intervention 309 children had one or more maltreatment-related event recorded in their records (adjusted mean rate per year 9.1 events per 1000 person years at risk; 95% CI 8.0–10.2; Table 2). In the year, following the intervention 226 children had one or more maltreatment-related event recorded (adjusted rate 12.3 events per 1000 person years at risk; 95% CI 10.7–14.2; Table 1). Rates of any maltreatment-related code increased by almost 40% (rate ratio: 1.4, 95% CI 1.1, 1.6; Table 3). The greatest

increase was seen in the ‘cause for concern’ code, especially for the older children (Table 3), although there were also increases in codes relating to child protection procedures following the intervention (Table 3).

### Process evaluation

Nine GP leads responded to the process evaluation questionnaire. Seven of these reported disseminating the guidance at a practice meeting. Other methods of dissemination included holding a specific training session, emailing guidance, distributing the information packs and through peer discussion. Six of the lead GPs reported that the majority of the GP within their practice were using the ‘Child is cause for concern’ read code following the intervention.

The simplicity of the coding strategy was cited as the most important factor in assisting implementation by five of

**Table 2** Number, person years and adjusted rates of coding before and after the simple coding intervention. Rates given are per 1000 person years at risk per calendar year. CI = confidence interval

	Denominator		Child with ≥1 maltreatment-related codes per calendar year		Child with a code reflecting a child protection procedure per calendar year		Child flagged ‘cause for concern’ per calendar year	
	Children (N)	Person years at risk	Children (n)	Adjusted rate (95% CI)	Children (n)	Adjusted rate (95% CI)	Children (n)	Adjusted rate (95% CI)
<b>Before intervention: 2010–2011</b>								
<b>Overall</b>	20,975	36,140	309	9.1 (8.0–10.2)	78	2.3 (1.8–3.0)	68	2.1 (1.6–2.8)
<b>&lt;5 years</b>	5697	9316	122	14.0 (11.4–17.2)	29	3.6 (2.3–5.5)	39	4.6 (3.2–6.6)
<b>5–18 years</b>	15,278	26,824	187	7.0 (6.0–8.0)	49	1.8 (1.3–2.4)	29	1.1 (0.7–1.6)
<b>After intervention: 2012</b>								
<b>Overall</b>	21,783	19,560	226	12.3 (10.7–14.2)	66	3.7 (2.8–4.8)	91	5.3 (4.2–6.7)
<b>&lt;5 years</b>	6468	5286	81	17.4 (13.5–22.3)	21	4.9 (2.9–8.0)	36	8.7 (5.8–12.5)
<b>5–18 years</b>	15,315	14,273	145	10.2 (8.6–12.0)	45	3.2 (2.3–4.2)	55	3.9 (2.9–5.0)

**Table 3** Age and gender profile adjusted rate ratios for recording child safeguarding codes after compared with before the simple coding intervention. CI = confidence interval

	Any child with ≥1 maltreatment-related codes	Any child with ≥1 code reflecting a child protection procedure	Any child coded as ‘cause for concern’
	Adjusted rate ratio (95% CI)	Adjusted rate ratio (95% CI)	Adjusted rate ratio (95% CI)
<b>Overall</b>	1.4 (1.1–1.6)	1.6 (1.1–2.2)	2.5 (1.8–3.4)
<b>&lt;5 years</b>	1.2 (0.9–1.6)	1.3 (0.8–2.3)	1.9 (1.2–2.9)
<b>5–18 years</b>	1.5 (1.2–1.8)	1.7 (1.1–2.6)	3.7 (2.4–5.9)

the GP leads. Time pressures in general practice, including lack of dedicated time for coding and changing IT systems, were cited as barriers to coding.

In teleconference feedback one GP lead felt education around coding safeguarding concerns had improved overall awareness of safeguarding. Other clinicians stated it was difficult to code this type of issue during the consultation because computer screens are generally visible to patients. There was clinician fear of damaging the patient–doctor relationship and fear of possible legal consequences if patients were aware of these codes being entered into the electronic record. Clinicians who did use these clinical codes stated they tended to code when patient or family had left the consultation room.

Some practices utilised safeguarding data entry templates to facilitate coding but found that free text entry was still used to record safeguarding concerns. Two of the practices reported that local policy and other local drivers meant that they had to use the code list selected by their locality. One local safeguarding board had imposed its own safeguarding template with different codes.

Administrative staff with specific responsibility for recording safeguarding data ‘Safeguarding Clerks’ already existed in one practice. They were responsible for coding data from clinical correspondence under the instruction of the GP clinical lead. They reported using the clinical codes listed. Another practice reported that coding in the consultation was left to the family physician, although a health visitor did have access to the notes and sometimes code cases in the practice computerised medical records. Other practices reported how health visitors are no longer based in the practice, and there was only sporadic sharing of information about children they considered to be a cause for concern.

## DISCUSSION

Despite time pressures and competing priorities within general practice, the simple coding strategy increased coding of maltreatment-related problems in children registered at 10 general practices in England. There were increases in codes relating to child protection procedures as well as in the ‘cause for concern’ code, which was the target of the coding strategy.

### Comparison with the literature

There is an estimated 4%–16% of children who are subject to abuse annually.<sup>5</sup> Despite improvements the coding rates seen here are much lower than this estimate. This gap may be explained by a continued reluctance by GPs to code maltreatment-related problems in the child’s records (14), sub-optimal recognition and/or sub-optimal use of health services use by maltreated and vulnerable children.

### Limitations of the method

The study is limited by the lack of a control group of practices. We are therefore unable to conclusively demonstrate that the improvements observed were caused by the

intervention and not by external factors, such as changes in safeguarding GP training, high profile safe guarding cases and other factors. Because the actual numbers of maltreatment-related codes in each practice are relatively small, the data are particularly sensitive to changes in case-mix within the practice. For example if a vulnerable family with several children were to register during the ‘after’ time period, that may result in a rise in maltreatment-related codes. However, such changes in case mix were unlikely to have occurred in all 10 practices and the increase that we saw exceeded the 10% per year increase which has been reported in analyses of large and representative sample of UK primary care data.<sup>2</sup>

Generalisability may be limited. GP practices involved were volunteer practices with GP leads that had an interest in child safeguarding or other relevant area of expertise who had been involved in the consensus development of the coding strategy. Implementation of this coding strategy in practices without the same buy-in may prove less effective. However, all UK practice are now required to have a child safeguarding lead, who could act as the ‘lead GP’ for implementation within their practice. Conversely, the impact of this coding strategy may be greater in non-expert practices as their baseline coding rate may be lower; three of the 10 practices in our sample already had coding strategies in place.

### Implications of the findings

There is no nationally agreed strategy for coding child maltreatment concerns in primary care. A national guideline would help guide local safeguarding boards and GP computer groups in each locality and facilitate improved recording. Application of the simple coding strategy improved coding of maltreatment-related problems and this approach could underpin any such national strategy. GPs can also adopt this coding strategy to improve their recording of maltreatment-related concerns.

### Further research

Continuity of care is important in establishing a therapeutic relationship in primary care and has many dimensions from good record keeping to personal continuity.<sup>15</sup> As the primary care workforce becomes more mobile, clinical record continuity becomes ever more important. Whilst national guidelines recommend coding the entire spectrum of maltreatment concerns<sup>8,9</sup> clinicians are very sensitive to the implications of putting codes into records that may affect their relationship with the patients.<sup>16</sup> Although there has been little analysis of whether GP concerns about recording safeguarding data are reflected in patient’s attitudes and beliefs, indirect evidence suggests that at least some parents are deterred from seeking care from a GP due to fears about their surveillance role and potential for a long-lasting and stigmatising label to be given to the child or family.<sup>17,18</sup> Further analysis is required into how acceptable it is to patient’s to have a ‘cause for concern’ code in their child’s record. This is particularly important as patients have

increasing access to their own records and especially as the UK's Department of Health has mandated that all family practices should offer online access to the records of their patients; effective in 2015.<sup>19</sup>

Given the policy emphasis from NICE guidance and GMC recommendations on recording, the use of read codes is a surrogate marker for the quality of child safeguarding in general practice. It is likely that recording of concerns is a necessary but not sufficient part of responding to child maltreatment and related problems. Whilst recording and coding concerns is the first step in responding we do not know how coding changes subsequent management of the child and family or what the impact of improving coding is on child or family outcomes. A quality improvement trial involving larger number of practices would enable exploration of whether care benefits for vulnerable children resulting from improving data recording outweigh the risks and concerns surrounding coding.

## CONCLUSION

A simple coding strategy can improve the coding of maltreatment-related concerns by GPs. Further research should investigate how recording can best support the doctor–patient relationship.

## REFERENCES

- Royal College of General Practitioners. Safeguarding Children & Young People: A Toolkit for General Practice 2011. Accessed: 03 Oct 2014. Available from: <http://www.rcgp.org.uk/~media/Files/CIRC/Safeguarding%20Children%20Module%20One/Safeguarding-Children-and-Young-People-Toolkit.ashx>.
- Woodman J, Freemantle N, Allister J, de Lusignan S, Gilbert R and Petersen I. Variation in recorded child maltreatment concerns in UK primary care records: a cohort study using The Health Improvement Network (THIN) database. *PLoS One* 2012;7(11):e49808. Epub 2012/12/05. <http://dx.doi.org/10.1371/journal.pone.0049808>. PMID:23209604; PMCID:PMC3509120.
- Woodman J, Allister J, Rafi I, de Lusignan S, Belsey J, Petersen I et al. A simple approach to improve recording of concerns about child maltreatment in primary care records: developing a quality improvement intervention. *The British journal of general practice: The Journal of the Royal College of General Practitioners* 2012;62(600):e478–86. Epub 2012/07/12. <http://dx.doi.org/10.3399/bjgp12X652346>. PMID:22781996; PMCID:PMC3381274.
- Finkelhor D, Turner H, Ormrod R and Hamby SL. Violence, abuse, and crime exposure in a national sample of children and youth. *Pediatrics* 2009;124(5):1411–23. Epub 2009/10/07. <http://dx.doi.org/10.1542/peds.2009-0467>. PMID:19805459.
- Gilbert R, Widom CS, Browne K, Fergusson D, Webb E and Janson S. Burden and consequences of child maltreatment in high-income countries. *Lancet* 2009;373(9657):68–81. Epub 2008/12/06. [http://dx.doi.org/10.1016/S0140-6736\(08\)61706-7](http://dx.doi.org/10.1016/S0140-6736(08)61706-7).
- Gonzalez-Izquierdo A, Woodman J, Copley L, van der Meulen J, Brandon M, Hodes D et al. Variation in recording of child maltreatment in administrative records of hospital admissions for injury in England, 1997-2009. *Archives of Disease in Childhood* 2010;95(11):918–25. Epub 2010/07/22. <http://dx.doi.org/10.1136/adc.2009.180216>. PMID:20647257.
- Woodman J, Brandon M, Bailey S, Belderson P, Sidebotham P and Gilbert R. Healthcare use by children fatally or seriously harmed by child maltreatment: analysis of a national case series 2005-2007. *Archives of Disease in Childhood* 2011;96(3):270–5. Epub 2011/01/19. <http://dx.doi.org/10.1136/adc.2010.190314>. PMID:21242231.
- National Institute for Health and Clinical Excellence. When to suspect child maltreatment (CG89)2009. Available from: <http://www.nice.org.uk/CG89>.
- General Medical Council. Protecting Children and Young People: The Responsibilities of All Doctors, 2012.
- Department for Education. Working together to safeguard children. 2013. Accessed: 03 Oct 2014. Available from: <https://www.gov.uk/government/publications/working-together-to-safeguard-children>.
- Kumarapeli P and de Lusignan S. Using the computer in the clinical consultation; setting the stage, reviewing, recording, and taking actions: multi-channel video study. *Journal of the American Medical Informatics Association* 2013;20(e1):e67–75. <http://dx.doi.org/10.1136/amiajnl-2012-001081>. Epub 2012/12/18.
- Royal College of General Practitioners. Safeguarding children and young people: a toolkit for general practice, 2011. Available

## NOTES

### Funding

The study of 11 practices was designed, registered and funded as an audit. The audit was coordinated by the RCGP-Clinical Innovation and Research Centre and funded by the Health Care Quality Improvement Partnership. Jenny Woodman was funded by a Medical Research Council/Economic and Social Research Council interdisciplinary Ph.D. studentship. Ruth Gilbert is an investigator with the Department of Health funded Policy Research Unit for Children, young people and families.

### Ethical considerations

This study observed the quality of recording of safeguarding concerns and the simple intervention was educating clinicians about how to implement a nationally agreed standard for recording. It is therefore defined as a clinical audit and as such did not require ethics committee review.<sup>20</sup> All practices consented to the extraction of anonymised data for audit purposes.

### Competing interests

The authors have declared no competing interests.

### Acknowledgements

The authors would like to thank the participating practices and GP leads.

from: <http://www.rcgp.org.uk/clinical-and-research/clinical-resources/child-and-adolescent-health/safeguarding-children-toolkit.aspx>.

13. de Lusignan S. Codes, classifications, terminologies and nomenclatures: definition, development and application in practice. *Informatics in Primary Care* 2005;13(1):65–70. Epub 2005/06/14: PMID:15949178.
14. Woodman J, Gilbert R, Allister J, Glaser D and Brandon M. Responses to concerns about child maltreatment: a qualitative study of GPs in England. *BMJ Open* 2013;3(12):e003894. Epub 2013/12/05. <http://dx.doi.org/10.1136/bmjopen-2013-003894>. PMID:24302506; PMCID:PMC3855649.
15. Haggerty JL, Roberge D, Freeman GK and Beaulieu C. Experienced continuity of care when patients see multiple clinicians: a qualitative metasummary. *Annals of Family Medicine* 2013;11(3):262–71. Epub 2013/05/22. <http://dx.doi.org/10.1370/afm.1499>. PMID:23690327; PMCID:PMC3659144.
16. de Lusignan S, Wells SE, Hague NJ and Thiru K. Managers see the problems associated with coding clinical data as a technical issue whilst clinicians also see cultural barriers. *Methods of Information in Medicine* 2003;42(4):416–22. Epub 2003/10/10. PMID:14534643.
17. Carvin K, Jones C, Marttila A, Burstrom B and Whitehead M. Can I risk using public services? Perceived consequences of seeking help and health care among households living in poverty: qualitative study. *Journal of Epidemiology and Community Health* 2007;61(11):984–9. Epub 2007/10/16. <http://dx.doi.org/10.1136/jech.2006.058404>. PMID:17933957; PMCID:PMC2465616.
18. Sayal K, Tischler V, Coope C, Robotham S, Ashworth M, Day C et al. Parental help-seeking in primary care for child and adolescent mental health concerns: qualitative study. *The British Journal of Psychiatry: The Journal of Mental Science* 2010;197(6):476–81. Epub 2010/12/02. <http://dx.doi.org/10.1192/bjp.bp.110.081448>. PMID:21119154.
19. de Lusignan S, Morris L, Hassey A and Rafi I. Giving patients online access to their records: opportunities, challenges, and scope for service transformation. *The British Journal of General Practice: The Journal of the Royal College of General Practitioners* 2013;63(611):286–7. Epub 2013/06/06. <http://dx.doi.org/10.3399/bjgp13X668032>. PMID:23735378, PMCID:PMC3662423.
20. National Research Ethics Service. Defining Research. 2009. Available from: <http://nuhriise.org/wp-content/uploads/Defining-Research-Leaflet.pdf>.